

INDUSTRIAL HEMP – FOR SEEDS, * FIBERS

FOLIAR FEEDING

Growth phase	Organogenetic phase	Use rates
Seedling emergence	I Differentiation of growth cone	1 l/ha
	II Differentiation of stem, initiation of leaves and axes of 2nd and subsequent orders	
Second pair of true leaves	III Differentiation of inflorescence axes	1 l/ha
	IV Initiation of flower organs	
	V Differentiation of flower organs	
	VI Formation of flower organs	
	VII Growth of flower organs	
Inflorescence formation	VIII Completion of organ formation in male and female flowers	1 l/ha
Flowering	IX Blooming and fertilization of female flowers, formation of bast fibers	1 l/ha
	X Formation of bast fibers, seeds	
Total per season		4 l/ha

Depending on the ripening performance of your crop variety, you may have to double the number of treatments within a phase.

The recommended product rates should be administered in a working solution applied at 200–300 l/ha. These application rates need not be changed when using low-volume and ultra-low-volume spraying techniques.

* We recommend to apply additional treatment at 1 l/ha after the topping.

MEDICAL HEMP

DRIP IRRIGATION

Growth phase	Germination	Seedling emergence	Vegetative	Flowering
How to use	—	Drip irrigation		
Use rates	—	0.05% solution based on the total amount of water is to be introduced into the feeding system on an ongoing basis		
				Stop feeding the plants with iGrow 2–3 weeks before harvest

ROOT ZONE WATERING

Growth phase	Germination	Seedling emergence	Vegetative	Flowering
How to use	—	Each watering		Stop feeding the plants with iGrow 2–3 weeks before harvest
Use rates	—	5 ml / 1 l of water		

FOLIAR FEEDING (SPRAYING)

Growth phase	Germination	Seedling emergence	Vegetative	Flowering	
How to use	—	Foliar application once in week starting from vegetative phase			Stop feeding the plants with iGrow 2–3 weeks before harvest
Use rates	—	2.5 ml / 1 l of water			

Recommended application method is in a tank mix combined with pesticides to de-stress plants.

Never add the product directly to the stock solution of the agrochemical.

Solution concentration 0,05% referenced to 10 l of water – 45 ml of iGrow B.

FOLIAR FEEDING

Our recommendations have been developed for professional applications based on data obtained in field tests. Consequently, all recommended rates are stated per tonne of seeds and per hectare of crop production area.

The recommended amount for foliar treatment should be administered in a working solution applied at the rate of 200–300 l/ha. The amount of fertilizer need not be changed when using low-volume and ultra-low-volume spraying techniques.

Winter and spring cereals**(Wheat, barley, oat, triticale, rye, millet)**

Tillering	0.3–0.5 l/ha
Booting	0.3–0.5 l/ha
Heading	0.3–0.5 l/ha
Milk stage	0.3–0.5 l/ha
Total per season	1.5–2.5 l/ha

Pulses**(Soy, pea, phaseolus, lentil, chickpea)**

Seedling emergence	0.3–0.5 l/ha
1–3 Leaves	0.3–0.5 l/ha
Branching	0.3–0.5 l/ha
Bud formation	0.3–0.5 l/ha
Flowering	0.3–0.5 l/ha
Total per season	1.5–2.5 l/ha

Maize

Seedling emergence	0.5–1.0 l/ha
3–5 Leaves	0.5–1.0 l/ha
6–8 Leaves	0.5–1.0 l/ha
Panicle emergence	0.5–1.0 l/ha
Total per season	2.0–4.0 l/ha

Sunflower

Seedling emergence	0.75–1.0 l/ha
1 Pair of leaves	0.75–1.0 l/ha
2–3 Pairs of leaves	0.75–1.0 l/ha
5–7 Pairs of leaves	0.75–1.0 l/ha
Bud formation	0.75–1.0 l/ha
Flowering	0.75–1.0 l/ha
Total per season	4.5–6.0 l/ha

Rape

Seedling emergence	0.3–1.0 l/ha
3–4 Leaves	0.3–1.0 l/ha
Rosette formation	0.3–1.0 l/ha
Stem elongation	0.3–1.0 l/ha
Bud formation	0.3–1.0 l/ha
Flowering	0.3–1.0 l/ha
Total per season	1.8–6.0 l/ha

Rice

Tillering	0.75–1.0 l/ha
Booting	0.75–1.0 l/ha
Panicle emergence	0.75–1.0 l/ha
Total per season	2.25–3.0 l/ha

Buckwheat, millet

3–5 Leaves	0.15–0.3 l/ha
Bud formation – Early flowering	0.15–0.3 l/ha
Flowering	0.15–0.3 l/ha
Total per season	0.45–0.9 l/ha

If your requirements are lower, just reduce the recommended amount by an appropriate percentage. For example, if your foliar application calls for 5 liters of our fertilizer per hectare of your potato field, then the per-acre equivalent will be as low as 2 liters, diluted in 120 liters of working solution.

Sugar beet

Seed leaves	0.3–1.0 l/ha
2 Leaves	0.3–1.0 l/ha
4 Leaves	0.3–1.0 l/ha
6 Leaves	0.3–1.0 l/ha
8 Leaves	0.3–1.0 l/ha
50 % Between-rows Crop cover	0.3–1.0 l/ha
Total per season	1.8–6.0 l/ha

Potato

Seedling emergence	0.75–1.25 l/ha
Stem elongation	0.75–1.25 l/ha
Bud formation	0.75–1.25 l/ha
Flowering	0.75–1.25 l/ha
Total per season	3.0–5.0 l/ha

Carrot

Seed leaves	0.3–1.0 l/ha
2 Leaves	0.3–1.0 l/ha
4 Leaves	0.3–1.0 l/ha
5–7 Leaves	0.3–1.0 l/ha
8–10 Leaves	0.3–1.0 l/ha
50 % Between-rows Crop cover	0.3–1.0 l/ha
Total per season	1.8–6.0 l/ha

Onion, garlic

Seedling emergence	0.25–0.75 l/ha
1–2 Leaves	0.25–0.75 l/ha
2–4 Leaves	0.25–0.75 l/ha
4–6 Leaves	0.25–0.75 l/ha
Vegetative mass growth	0.25–0.75 l/ha
Bulb formation	0.25–0.75 l/ha
Total per season	1.5–4.5 l/ha

CABBAGE

Seedling emergence	0.75–1.0 l/ha
1–2 Leaves	0.75–1.0 l/ha
2–4 Leaves	0.75–1.0 l/ha
Early head formation	0.75–1.0 l/ha
Head fill	0.75–1.0 l/ha
Total per season	3.75–5.0 l/ha

Tomato

Seedling emergence	0.5–1.0 l/ha
1–2 Leaves	0.5–1.0 l/ha
2–4 Leaves	0.5–1.0 l/ha
6–8 Leaves	0.5–1.0 l/ha
Bud formation	0.5–1.0 l/ha
Flowering	0.5–1.0 l/ha
Fruit set	0.5–1.0 l/ha
Total per season	3.5–7.0 l/ha

Bell pepper, eggplant

Seedling emergence	0.5–1.0 l/ha
1–2 Leaves	0.5–1.0 l/ha
2–4 Leaves	0.5–1.0 l/ha
3–5 Leaves	0.5–1.0 l/ha
Bud formation	0.5–1.0 l/ha
Flowering	0.5–1.0 l/ha
Fruit set	0.5–1.0 l/ha
Total per season	3.5–7.0 l/ha

Cucumber, zucchini

Seedling emergence	0.5–1.0 l/ha
1–2 Leaves	0.5–1.0 l/ha
2–4 Leaves	0.5–1.0 l/ha
6–8 Leaves	0.5–1.0 l/ha
Bud formation	0.5–1.0 l/ha
Flowering	0.5–1.0 l/ha
Fruit set	0.5–1.0 l/ha
Total per season	3.5–7.0 l/ha

Melon

Seedling emergence	0.5–1.0 l/ha
1–2 Leaves	0.5–1.0 l/ha
2–4 Leaves	0.5–1.0 l/ha
6–8 Leaves	0.5–1.0 l/ha
Bud formation	0.5–1.0 l/ha
Flowering	0.5–1.0 l/ha
Fruit set	0.5–1.0 l/ha
Total per season	3.5–7.0 l/ha

Garden strawberry

Springtime leaf regrowth	0.3–0.75 l/ha
Early flower-bud formation	0.3–0.75 l/ha
Flowering	0.3–0.75 l/ha
Post-harvest care	0.3–0.75 l/ha
Total per season	1.2–4.0 l/ha

Grape

Budbreak	0.75–1.5 l/ha
2–5 Leaves	0.75–1.5 l/ha
5–8 Leaves	0.75–1.5 l/ha
Flower cluster initiation, flowering	0.75–1.5 l/ha
Late flowering, fruit set	0.75–1.5 l/ha
Berry growth	0.75–1.5 l/ha
Berry coloring	0.75–1.5 l/ha
Total per season	5.25–10.5 l/ha

Pip fruits

'Green cone'	0.75–2.25 l/ha
Flower-bud separation	0.75–2.25 l/ha
'Pink bud'	0.75–2.25 l/ha
'Hazelnut'	0.75–2.25 l/ha
'Walnut'	0.75–2.25 l/ha
Fruit growth	0.75–2.25 l/ha
Total per season	4.5–13.5 l/ha

Stone fruits

Late flowering	0.75–2.25 l/ha
Formation of petiolar fossa	0.75–2.25 l/ha
Fruit growth	0.75–2.25 l/ha
Post-harvest care	0.75–2.25 l/ha
Total per season	2.0–9.0 l/ha

Lawn grasses

Early regrowth	50 ml / 10 l of water
Vegetation	50 ml / 10 l of water
Late vegetation	50 ml / 10 l of water

Flowers

Bulbs, cuttings	50 ml / 10 l of water
Early vegetation	50 ml / 10 l of water
Bud formation	50 ml / 10 l of water
Early flowering	50 ml / 10 l of water

Ornamental shrubs

Cuttings, transplants	50 ml / 10 l of water
Budbreak	50 ml / 10 l of water
Bud formation	50 ml / 10 l of water
Flowering	50 ml / 10 l of water
Vegetative mass growth	50 ml / 10 l of water